ST030

GL Sciences Inc.


Analysis of Cyromazine in Livestock Products

Cyromazine is triazine based molecule that is used as an insect growth inhibitor for the protection of livestock and poultry houses. Standard values have been set for its use in veterinary drugs, and a notification method using HPLC has been issued.

In this study, extraction and purification were performed with a mixed-mode polymer SPE (InertSep MCX FF) using a method based on the notification test method from the Ministry of Health, Labor and Welfare and literature examples*, and analysis was performed using HPLC (detection at UV 240nm) and LC-MS/MS.

*Tokyo Metropolitan Institute of Public Heath Annual Report No.57 (2006)

SPE Pretreatment Flow



[InertSep MCX FF]



InertSep MCX FF is a styrene-divinylbenzene polymer (SDB) solid phase modified with a strong cation exchange group.

By exerting both reverse phase and cation exchange actions, basic compounds can be strongly retained.

onal group strong

Since the particle size is up to 70 μ m, it is ideal for biological sample extracts.

Recovery test with spiked extract (milk, 0.05 mg / kg, n = 2)


 H_2N


Cyromazine
Structures created using Chemistry 4-D Draw provided by ChemInnovation Software, Inc.

Rec (%)	HPLC-PDA	LC-MS/MS
Milk	80.1	78.6

Average Particle Size: 70 µm Pore Volume: 1.1 mL/g pH Range: 1-14 Surface Area: 480 m2/g Pore diameter: 9nm

Measurement conditions





HPLC conditions

〈HPLC-PDA〉

Column : InertSustain C18

 $(3 \mu m, 150 \times 4.6 \text{ mm I.D.})$

Column Cat. No. : 5020-07445

Eluent : A) 50 mmol/L CH₃COONH₄

B) CH₃CN A/B=49/1, v/v

Flow rate : 1.0 mL/min
Col. Temp. : 40 °C
Detection : PDA 240 nm
Injection Vol. : 50 μL

Sample : 1. Cyromazine 1.75 mg/L

(LC-MS/MS)

Column : InertSustain C18

(3 μm, 150 x 2.1 mm I.D.)

Column Cat. No. : 5020-14355

Eluent : A) 50 mmol/L CH₃COONH₄

B) CH_3CN A/B=49/1, v/v

Flow rate : 0.4 mL/min
Col. Temp. : 40 °C
Detection : LC/MS/MS

(4000QTRAP: ESI, Positive, SRM)

Injection Vol. : 10 μL

Sample : 1. Cyromazine 1.75 mg/L

Q1 / Q3 = 167.0 / 68.0 (Quantifier ion)

167.0 / 85.0 (Qualifier ion)

3. Applicability to other samples


0.5 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0 5.5 6.0

Time, min


Depending on the sample matrix, sufficient recovery of the target components may not be possible. If the matrix affects recovery, increasing the loading amount on the MCX cartridge may improve the recovery rate. However, if the loading amount is increased, peaks from interfering components will also increase, so it is important to select an appropriate loading amount suitable for the matrix.

6.5 7.0 7.5 8.0 8.5 9.0 9.5

<Other samples: chicken>



Recovery rate 97.5%



GL Sciences disclaims any and all responsibility for any injury or damage which may be caused by this data directly or indirectly. We reserve the right to amend this information or data at any time and without any prior announcement.

GL Sciences Inc. Japan

22-1 Nishishinjuku 6-chome Shinjuku-ku, Tokyo 163-1130, Japan

Phone: +81-3-5323-6620 Fax: +81-3-5323-6621 Email: <u>world@gls.co.jp</u> Web: www.glsciences.com

GL Sciences Inc. USA

4733 Torrance Blvd. Suite 255 Torrance, CA 90503 USA

Phone: +1-310-265-4424
Fax: +1-310-265-4425
Email: info@glsciencesinc.com
Web: www.glsciencesinc.com

GL Sciences B.V.

Dillenburgstraat 7C 5652AM, Eindhoven The Netherlands

Phone: +31-40-254-9531 Email: info@glsciences.eu Web: www.glsciences.eu

GL Sciences (Shanghai) Limited

Tower B, Room 2003 Far East International Plaza No.317 Xianxia Road, Changning District Shanghai, China 200051

Phone: +86-21-62782272

Email: contact@glsciences.com.cn Web: www.glsciences.com.cn



Visit our Website at www.glsciences.com/distributors